

The Gyrokinetic Regime

Geometry

Velocity Space

Linear How-To Units

Non-Linear Issues

Miscellany

Coordinates

- Recall GS2 uses energy E and pitch angle $\lambda = \mu/E$
- λ grid is divided into trapped and untrapped regions
- Untrapped pitch angle resolution determined by `ngauss`
- Select `ngauss` from this list:

1, 2, 3, 4, 5, 6, 8, 10, 12, 16, 20, 24, 32, 40, 48

Why Choices are Restricted

- E, λ grids used in collision operator & source integrals
- The energy grid and the untrapped component of the pitch angle grid are pseudo-spectral to improve accuracy
- Simpler rules like

$$\int_a^b f(x) dx = \sum_{i=1}^N f(x_i) w_i$$

integrate polynomial of degree $N + 1$ accurately over finite domain with equally spaced grid points.

Pseudo-spectral grids (Gaussian quadrature)

- But, use freedom to choose x_i to minimize error; increases degree of polynomials that are exactly integrated to $2N + 1$.
- No free lunch: x_i, w_i must be precomputed.
- Precomputed untrapped pitch angle grids hardwired.
- Upgrade expected soon, following Candy & Waltz
- Energy grid already upgraded. . .

Energy Grid

- Two choices for energy grid discretization available
- Namelist `le_grids_knobs`, `advanced_egrid = T` selects the Candy-Waltz algorithm and is the default. With this choice:
 1. Set `negrid > 1` (any integer)
 2. Set `ecut ~ 6.0`
- Otherwise (to compare with old results, mainly), consult website for guidance.

Energy Grid: Note for Developers

- Internally, the energy grid is normalized according to the definition $v_t \equiv \sqrt{2T/m}$ regardless of any choices in the input file.
- It is possible to turn off trapped particles, but this option is not fully supported yet and may not work.

Collision Operator

- Lorentz collision operator:

$$C(f) = \frac{\nu(E)}{2} \frac{\partial}{\partial \xi} (1 - \xi^2) \frac{\partial f}{\partial \xi}$$

with optional momentum-conserving terms allowed (but not recommended!)

- For example, for electron-ion collisions:

$$\nu(E) = \frac{\nu_{ei}}{(v/v_{te})^3} \left[Z_{\text{eff}} + \frac{v_{te}}{v\sqrt{\pi}} e^{-(\frac{v}{v_{te}})^2} + \left(1 - \frac{v_{te}}{2v^2}\right) \text{erf}\left(\frac{v}{v_{te}}\right) \right]$$

where

$$\nu_{ei} = \frac{4\pi n e^4 \ln \Lambda}{(2T_e)^{3/2} m_e^{1/2}}$$

Collision Operator Inputs

- Recommend `collision_model = "default"`, (automatically selected if choice is left out of input file)
- Recommend `conserve_momentum = F`. The physics model isn't reliable, and the performance takes a big hit.
- Set collision frequency for each species in `species_knobs` namelist (variable `vnewk`)
- See detailed notes on website:
http://gs2.sourceforge.net/collision_notes.ps

Adiabaticity

- Sometimes it is wise to make one species adiabatic:
 1. For ITG studies, might wish to use adiabatic electrons
 2. For ETG studies, might wish to use adiabatic ions
- If no species has `type = 'electron'` an adiabatic species is `automatically` added.
- This feature can be used to simulate ETG modes with adiabatic ions, by pretending the electrons are ions with charge $Z = -1$

Adiabaticity Options

- The form of the adiabatic response can be chosen from:
 1. Namelist `dist_fn_knobs`, `adiabatic_option='default'`
 - Appropriate for adiabatic ions in an ETG simulation
 2. Namelist `dist_fn_knobs`, `adiabatic_option='field-line-average-term'`
 - Appropriate for adiabatic electrons in an ITG simulation

Beams and Impurities

- Okay to leave out charge from such species.
- Beams can be mocked up with high-temperature Maxwellian